

Equivariant deformations of the affine multicone over a flag variety

P. Bravi and S. Cupit-Foutou

Abstract

We prove that the invariant Hilbert scheme parametrising the equivariant deformations of the affine multicone over a flag variety is, under certain hypotheses, an affine space. The proof is based on the construction of a wonderful variety in a fixed multiprojective space.

Introduction

Given a finite dimensional G -module V with G a complex connected reductive group, the invariant Hilbert scheme parameterises the closed G -subschemes of V whose coordinate ring regarded as a G -module is isomorphic to a fixed G -module. This fixed G -module defines an invariant – the Hilbert function – which is the analog of the Hilbert polynomial in case of Grothendieck’s Hilbert scheme. In [AB], V. Alexeev and M. Brion studied the invariant Hilbert scheme and proved that it was quasiprojective.

Of special interest ([AB]) are the invariant Hilbert schemes whose corresponding Hilbert function takes 0 or 1 as only values, namely the fixed G -module is multiplicity-free. Their closed points are the so called affine spherical varieties (e.g. affine toric varieties).

The coordinate ring of the affine cone of primitive vectors in a simple G -module V is multiplicity-free. The equivariant deformations of such a cone are parameterised by the invariant Hilbert scheme for the corresponding Hilbert function. In [J], S. Jansou proved that this invariant Hilbert scheme was a (reduced) point or an affine line.

In this paper we consider the more general situation given by an affine multicone over a flag variety, provided it is normal, spherical and with boundary of codimension at least two.

These hypotheses we make on the Hilbert function are not new in invariant deformation theory (see [Pa] and also [VP, BK, Bri86, K90]). They

yield a correspondence between deformations of spherical G -varieties and deformations of their open G -orbits.

Under these assumptions, we thus prove that the invariant Hilbert scheme, parameterising the equivariant deformations of an affine multicone, is an affine space. Its dimension can be explicitly determined: it is closely related to a combinatorial datum (called spherical system) that we can attach to the Hilbert function.

In [J], the cases where the Hilbert scheme is the affine line are explicitly listed. This list corresponds to that of the two-orbit varieties whose closed orbit is a very ample divisor ([A] and [HS, Bri89]).

The two-orbit varieties which appear in the above picture are wonderful varieties. The wonderful varieties are spherical ([L96]), the knowledge of this special class of varieties should lead to a better understanding of spherical varieties (*see* [L01]). In general, to any wonderful variety one can associate a spherical system, but the converse remains an open problem; only partial results are known (*loc. cit.* and [Pe03, BP, Bra]).

This paper is an attempt to clarify the connection between invariant Hilbert schemes and wonderful varieties.

In this setting we prove that the total cone of deformations of the multicone over the flag variety we consider is given by the multicone over a well determined wonderful G -variety containing that flag variety as its unique closed G -orbit.

In the first section we recall the definition of the invariant Hilbert scheme in the multiplicity free case. We focus on its closed points and define the action of the adjoint torus. We also briefly recall some definitions concerning wonderful varieties.

In the second section we introduce our affine multicone over a flag variety under certain assumptions and we state our main results.

The remaining sections are dedicated to the proofs.

By means of a representation theoretical characterisation of the tangent space of the invariant Hilbert scheme at the multicone given in [AB], we compute explicitly in the third section the weights of the toric action on this tangent space. We prove also that these weights are multiplicity free.

In the fourth section we recall the definition of spherical systems, due to D. Luna. We then associate a spherical system to the considered invariant Hilbert function.

Afterward, we show that a wonderful spherical subgroup N corresponding to such a spherical system provides an open orbit in the Hilbert scheme. Furthermore, by means of a result of G. Pezzini, we prove that the compactification of G/N in the given multiprojective space is the investigated wonderful variety.

In the last section we show how to associate to a spherical system in our setting a corresponding wonderful subgroup.

Acknowledgements. This work is mostly the result of a staying of the first author at the Bergische Universität Wuppertal while supported by an European “Marie Curie” TMR *Liegrits* fellowship. Both authors would like to thank M. Brion, S. Jansou, P. Littelmann and D. Luna for helpful discussions, suggestions and support.

Notation. The ground field is the field of complex numbers. Given a connected, simply connected, semisimple algebraic group G and a maximal torus T , let Λ be the lattice of T -weights and Φ its root system. Fixing a Borel subgroup B containing T , we denote by $S = \{\alpha_1, \dots, \alpha_n\}$ the associated set of simple roots and by Φ^+ the set of positive roots. To each root β , we attach a root vector X_β of the Lie algebra $\mathfrak{g}$ of G .

Let Λ^+ be the cone of dominant weights. Recall that $\Lambda^+ = \bigoplus_1^n \mathbb{N}\omega_i$ where ω_i is the fundamental weight attached to α_i . If $\alpha_j^\vee$ is the coroot of α_j , we have $(\omega_i, \alpha_j^\vee)$ equal to 1 if $i = j$ and equal to 0 otherwise.

The simple G -modules are in correspondence with the cone Λ^+ of dominant weights. We thus denote by $V(\lambda)$ the simple G -module with highest weight $\lambda \in \Lambda^+$.

1 Definitions

1.1 Invariant Hilbert scheme

Let Γ be a submonoid of Λ^+ generated by dominant weights $\lambda_1, \dots, \lambda_s$. Denote by V the finite dimensional G -module given by $\bigoplus_{i=1}^s V(\lambda_i)$, and by R the G -module given by $\bigoplus_{\lambda \in \Gamma} V(\lambda)^*$.

In this setting, we follow [AB] to define (and state some properties of) the *invariant Hilbert scheme* $\text{Hilb}_\Gamma^G(V)$ of affine G -subschemes of V with G -module of regular functions isomorphic to R . In a more general context, the invariant Hilbert scheme is defined up to the *Hilbert function* instead of the weight monoid Γ . In the multiplicity free case, these two data are equivalent so we will not recall here the definition of this function.

A *family* of affine G -subschemes of V over a given scheme S is a closed G -subscheme $\mathfrak{X} \subset V \times S$ where the restriction on $\mathfrak{X}$ of the projection $\pi: V \times S \rightarrow S$ is affine and of finite type.

Let $\mathcal{R}$ denote the direct image $\pi_* \mathcal{O}_{\mathfrak{X}}$ of the structure sheaf of $\mathfrak{X}$. Such a family is said to be *of type* Γ if we have

$$\mathcal{R} \cong \bigoplus_{\lambda \in \Gamma} \mathcal{R}_\lambda \otimes V(\lambda)^* \quad (1.1)$$

and each $\mathcal{R}_\lambda$ is an invertible sheaf of $\mathcal{O}_S$ -modules. In particular the map $\pi: \mathfrak{X} \rightarrow S$ is flat.

Theorem 1.1 ([AB] Theorem 1.7). *The contravariant functor $\text{Hilb}_\Gamma^G(V): \{\text{schemes}\} \rightarrow \{\text{sets}\}$, that to any scheme S associates the set of families $\mathfrak{X} \subset V \times S$ of affine G -subschemes of V of type Γ , is represented by a quasi-projective scheme.*

The invariant Hilbert scheme $\text{Hilb}_\Gamma^G(V)$ is such a quasi-projective scheme.

1.2 Closed points

An algebraic variety endowed with an action of G is said to be *spherical* if it is normal and it contains an open orbit of a Borel subgroup of G .

An affine G -variety X is spherical if and only if its algebra of regular functions $\mathbb{C}[X]$ is a *multiplicity-free* G -module, namely it is the direct sum of simple modules each of them occurring with multiplicity one. In this case, let $\Gamma \subset \Lambda^+$ be the monoid such that

$$\mathbb{C}[X] \cong \bigoplus_{\lambda \in \Gamma} V(\lambda)^* \quad (1.2)$$

as G -modules. The normality of X implies $\mathbb{Z}\Gamma \cap \mathbb{Q}_{\leq 0}\Gamma = \Gamma$.

A spherical G -subvariety of V with weight monoid equal to Γ can be regarded as a closed point of $\text{Hilb}_\Gamma^G(V)$. It is said to be *non-degenerate* if all its projections into the isotypical components of V are non-zero.

Theorem 1.2 ([AB] Corollary 1.17). *The non-degenerate spherical G -subvarieties of V with weight monoid Γ are parameterised by an open subscheme $\text{Hilb}_\Gamma^G(V)_0$ of $\text{Hilb}_\Gamma^G(V)$.*

Let X_0 and X_1 be two quasi-affine spherical G -varieties, X_0 is said to be a *contraction*, or a *degeneration* of X_1 , or vice versa that X_1 is a *deformation* of X_0 , if $\mathbb{C}[X_0]$ is G -isomorphic to the graduate algebra associated to a filtration of $\mathbb{C}[X_1]$.

Note that on a G -algebra $R \cong \bigoplus_{\lambda \in \Gamma} V(\lambda)^*$ we have the maximal filtration $\mathfrak{F}_\mu = \bigoplus_{\lambda \leq \mu} V(\lambda)^*$, for all $\mu \in \Gamma$, where $\lambda \leq \mu$ means that $\mu - \lambda$ is a sum of positive roots.

1.3 Toric action on the invariant Hilbert scheme

Let $Z(G)$ be the center of G . The invariant Hilbert scheme $\text{Hilb}_\Gamma^G(V)$ is endowed with an action of the adjoint torus $T_{\text{ad}} = T/Z(G)$ (see [AB]). The

open subscheme $\text{Hilb}_\Gamma^G(V)_0$ is stable for this action. This action can be defined on multiplication laws of $R = \bigoplus_{\lambda \in \Gamma} V(\lambda)^*$ as follows.

Let X be a non-degenerate spherical G -subvariety of V with weight monoid Γ . Its G -algebra of regular functions is isomorphic to R defined above, as G -module. This gives a G -multiplication law in R , $m: R \otimes R \rightarrow R$. The map m is direct sum of maps $m_{\lambda,\mu}^\nu: V(\lambda)^* \otimes V(\mu)^* \rightarrow V(\nu)^*$ where λ, μ, ν are in Γ , with $\nu \leq \lambda + \mu$.

For all $t \in T$, set

$$t.m = \bigoplus_{\lambda,\mu,\nu} t^{\lambda+\mu-\nu} m_{\lambda,\mu}^\nu. \quad (1.3)$$

The limit of $t.m$ for t^α going to zero, for some $\alpha \in S$, gives rise to a degeneration of m . If t^α goes to zero, for all $\alpha \in S$, we obtain the so called *Cartan multiplication*, that is $m_{\lambda,\mu}^\nu$ nonzero only if $\nu = \lambda + \mu$, corresponding to the maximal degeneration.

1.4 Wonderful varieties

An algebraic G -variety X is said to be *wonderful of rank r* if

- (i) X is smooth and complete
- (ii) G has an open G -orbit whose complement is the union of smooth prime G -divisors D_i ($i = 1, \dots, r$) with normal crossings such that $\bigcap_1^r D_i \neq \emptyset$
- (iii) if x, x' are such that $\{i : x \in D_i\} = \{i : x' \in D_i\}$ then $G.x = G.x'$.

A wonderful variety X is projective and spherical (see [L96]).

The *spherical roots* of X are the T -weights in the quotient $T_z X / T_z Y$, where Y is the unique closed G -orbit of X and $z \in Y$ is the unique point fixed by the opposite Borel subgroup of B . The rank of X is equal to the number of spherical roots of X .

Note that wonderful varieties of rank 0 are homogeneous: they are the flag varieties.

2 Main results

2.1 The saturation assumption

Definition 2.1 ([Pa]). A submonoid of dominant weights, $\Gamma \subseteq \Lambda^+$, is said to be *saturated* if

$$\mathbb{Z}\Gamma \cap \Lambda^+ = \Gamma. \quad (2.1)$$

The following characterisation of saturation for a free monoid is straightforward.

Lemma 2.1. *Let $\lambda_1, \dots, \lambda_s$ be linearly independent dominant weights. The semigroup $\Gamma = \langle \lambda_1, \dots, \lambda_s \rangle_{\mathbb{N}}$ is saturated if and only if for every $i = 1, \dots, s$ there exists a simple root α_{k_i} which is not orthogonal only to the weight λ_i .*

2.2 The affine multicone over a flag variety

Let $\lambda_1, \dots, \lambda_s$ be dominant weights. For all i , denote P_{λ_i} the stabiliser of the line $[v_{\lambda_i}] \in \mathbb{P}(V(\lambda_i))$ generated by a highest weight vector $v_{\lambda_i} \in V(\lambda_i)$. Let $P = \cap_1^s P_{\lambda_i}$ that is, the stabiliser of $([v_{\lambda_1}], \dots, [v_{\lambda_s}])$ in $\mathbb{P}(V(\lambda_1)) \times \dots \times \mathbb{P}(V(\lambda_s))$. We have the following obvious inclusions

$$G/P \subset G/P_{\lambda_1} \times \dots \times G/P_{\lambda_s} \subset \mathbb{P}(V(\lambda_1)) \times \dots \times \mathbb{P}(V(\lambda_s)). \quad (2.2)$$

Let X_0 denote the corresponding *affine multicone* over the partial flag variety G/P . It is the affine G -subvariety of $V = \oplus_{i=1}^s V(\lambda_i)$ defined as

$$X_0 = \{v_1 + \dots + v_s \in V : \exists ([u_1], \dots, [u_s]) \in G/P \text{ with } v_i \in [u_i] \forall i\}. \quad (2.3)$$

It is a subvariety contained in the product $\mathcal{C}_{\lambda_1} \times \dots \times \mathcal{C}_{\lambda_s}$ of the affine cones $\mathcal{C}_{\lambda_i} \subset V(\lambda_i)$ over $G/P_{\lambda_i} \subset \mathbb{P}(V(\lambda_i))$.

If the dominant weights $\lambda_1, \dots, \lambda_s$ are supposed to be linearly independent then the G -orbits of X_0 are parameterised by the subsets I of $\{1, \dots, s\}$,

$$G.v_{\underline{\lambda}}^I = \{v_1 + \dots + v_s \in X_0 : v_i = 0 \forall i \in I\}, \quad (2.4)$$

where $v_{\underline{\lambda}}^I = \sum_{i \notin I} v_{\lambda_i}$. In particular, the G -orbit of $v_{\underline{\lambda}} = (v_{\lambda_1}, \dots, v_{\lambda_s})$ is dense in X_0 .

The G -variety X_0 is normal with an open G -orbit. Further, its generic stabiliser is *horospherical*, i.e. it contains a maximal unipotent subgroup. In particular, the G -variety X_0 is spherical.

Denote Γ the linear span of $\lambda_1, \dots, \lambda_s$ over $\mathbb{N}$ and, moreover, assume it to be saturated. By Lemma 2.1 we get that, for any nonempty subset I , the stabiliser $G_{v_{\underline{\lambda}}}$ is included in $G_{v_{\underline{\lambda}}^I}$ with codimension at least two, therefore

$$\text{codim}_{X_0}(X_0 \setminus G.v_{\underline{\lambda}}) \geq 2. \quad (2.5)$$

The ring of regular functions $\mathbb{C}[X_0]$ is thus equal to $\mathbb{C}[G/G_{v_{\underline{\lambda}}}]$, hence isomorphic to $\oplus_{\lambda \in \Gamma} V(\lambda)^*$ as a G -module.

The non-degenerate spherical G -subvariety X_0 of V corresponds to the unique T_{ad} -stable closed point in $\text{Hilb}_{\Gamma}^G(V)_0$ and any non-degenerate spherical G -subvariety of V with weight monoid Γ is a deformation of X_0 .

2.3 Statements

From now on, $\lambda_1, \dots, \lambda_s$ are linearly independent dominant weights generating a saturated monoid Γ and V is the G -module given by $V(\lambda_1) \oplus \dots \oplus V(\lambda_s)$. The parabolic subgroup P is the stabiliser of $([v_{\lambda_1}], \dots, [v_{\lambda_s}])$ in $\mathbb{P}(V(\lambda_1)) \times \dots \times \mathbb{P}(V(\lambda_s))$, X_0 is the affine multicone over G/P included in this multi-projective space, as defined above.

Theorem 2.2. *As a T_{ad} -module the tangent space of $\text{Hilb}_{\Gamma}^G(V)$ at the point X_0 is multiplicity free and its T_{ad} -weights are spherical roots.*

In the following we shall denote the above set of T_{ad} -weights by Σ .

Theorem 2.3. (i) *There exists a wonderful G -variety X included in the multiprojective space $\mathbb{P}(V(\lambda_1)) \times \dots \times \mathbb{P}(V(\lambda_s))$ with set of spherical roots Σ and with closed G -orbit G/P .*

(ii) *Let $\tilde{X}$ be the corresponding affine multicone over X . Then there exists $v = (v_1, \dots, v_s) \in \tilde{X}$, with $([v_1], \dots, [v_s])$ in the open G -orbit of X , such that the subvariety $X_1 = \overline{G.v}$ of V is a deformation of X_0 .*

Corollary 2.4. *The invariant Hilbert scheme $\text{Hilb}_{\Gamma}^G(V)_0$ of non-degenerate spherical G -subvarieties of V with weight monoid Γ is an affine space. Its dimension is equal to the rank of the wonderful variety X of Theorem 2.3.*

In particular, the dimension of the invariant Hilbert scheme is less or equal than s , the rank of the free monoid Γ . Moreover, the invariant Hilbert scheme is a point if and only if the wonderful variety X given above is of rank 0, namely $X = G/P$.

In Corollary 5.5 the cases where the invariant Hilbert scheme $\text{Hilb}_{\Gamma}^G(V)_0$ is not a point will be characterised by an explicit condition on the monoid Γ .

Proof. The spherical subvariety X_1 of V given in Theorem 2.3 corresponds to a closed point in $\text{Hilb}_{\Gamma}^G(V)_0$. We shall prove that the T_{ad} -orbit of X_1 in $\text{Hilb}_{\Gamma}^G(V)_0$ is dense.

By Corollary 2.14 in [AB] the normalisation of the closure of the T_{ad} -orbit of X_1 in $\text{Hilb}_{\Gamma}^G(V)_0$ is isomorphic to a T_{ad} -module whose T_{ad} -weights are explicitly given. They are the linear independent generators of the free submonoid

$$\mathbf{M} = \mathbb{Q}_{\geq 0} \mathbf{M}' \cap \mathbb{Z} \mathbf{M}' \subseteq \mathbf{N} S,$$

obtained from the submonoid

$$\mathbf{M}' = \langle \lambda + \mu - \nu : \lambda, \mu, \nu \in \Gamma, m_{\lambda, \mu}^{\nu} \neq 0 \rangle_{\mathbb{N}},$$

where $m_{\lambda,\mu}^\nu: V(\lambda)^* \otimes V(\mu)^* \rightarrow V(\nu)^*$ and $m = \sum m_{\lambda,\mu}^\nu$ is the multiplication in $\mathbb{C}[X_1]$ (see Paragraph 1.3).

From Theorem 1.3 in [K96] we know that such T_{ad} -weights are exactly the spherical roots of the spherical variety X_1 , which are in correspondence (but maybe not equal) to the spherical roots in Σ . Therefore, by Theorem 2.2, the dimension of the orbit $T_{\text{ad}}.X_1$ is equal to the dimension of the tangent space $T_{X_0}\text{Hilb}_\Gamma^G(V)$. It follows that $\text{Hilb}_\Gamma^G(V)_0$ is smooth, hence equal to $\overline{T_{\text{ad}}.X_1}$ and isomorphic to an affine space. $\square$

Remark 2.1. Let us keep the notation of the previous proof. By Proposition 2.13 in [AB] the monoid $\mathbf{M}'$ is the weight monoid of $\overline{T_{\text{ad}}.X_1}$ in $\text{Hilb}_\Gamma^G(V)$. Therefore we have, a posteriori, that $\mathbf{M}'$ is free.

Corollary 2.5. *The union of all non-degenerate spherical G -subvarieties of V with weight monoid Γ is equal to the affine multicone $\tilde{X}$ over the wonderful G -variety X of Theorem 2.3 included in $\mathbb{P}(V(\lambda_1)) \times \dots \times \mathbb{P}(V(\lambda_s))$.*

The union of all non-degenerate spherical G -subvarieties of V with weight monoid Γ is a closed subset of V . By the same arguments as in the proof of Proposition 5.1 one can get that for any point $([v'_1], \dots, [v'_s])$ in X , $v' = (v'_1, \dots, v'_s)$ is such that the subvariety $\overline{G.v'}$ of V is a deformation of X_0 . Then the above corollary will follow from Theorem 2.3(i) and Corollary 2.4.

2.4 The case of a simple module

As already mentioned in the introduction, S. Jansou considers in [J] the case where the monoid Γ is generated by a single dominant weight λ and proves that the corresponding Hilbert scheme is a point or an affine line. Moreover, he discovers a correspondence with the list of one dimensional Hilbert schemes and the list of two-orbit varieties whose closed orbit is an ample divisor.

The two-orbit varieties X' which appear in this setting are (by definition) wonderful of rank one.

The union of the equivariant deformations of X_0 (here X_0 is the cone of the primitive vectors in $V(\lambda)$) is obtained by taking the affine cone over the wonderful variety X (still of rank one) whose generic stabiliser is the normaliser of the generic stabiliser of the former rank one wonderful variety X' .

2.5 The model case

On the other side D. Luna in [L05] considers the case where the monoid Γ is equal to Λ^+ . More precisely he is concerned with the so called *model ho-*

mogenous spaces, that are the quasi-affine (spherical) G -homogeneous spaces whose ring of regular functions is isomorphic to $\oplus_{\lambda \in \Lambda^+} V(\lambda)^*$ as a G -module. He obtains that their isomorphism classes are parameterised by the G -orbits of a certain wonderful variety for a given connected reductive group G . Further, this wonderful variety can be constructed explicitly case by case.

2.6 On the saturation assumption

Let H be a spherical subgroup of G , that is the homogeneous space G/H has an open B -orbit. The *horospherical contraction* $\widehat{H}$ of H ([Pa], [K90]), in the case of a quasi-affine G/H , is defined as follows.

Let $\Gamma(G/H)$ be the submonoid of highest weights in $\mathbb{C}[G/H]$, then $\widehat{H}$ is the intersection of the stabilisers G_{v_λ} for all weights λ in $\Gamma(G/H)$.

The subgroup $\widehat{H}$ is horospherical. Moreover, $\dim G/\widehat{H} = \dim G/H$ and ([Pa] Proposition 1.5)

$$\Gamma(G/\widehat{H}) = \mathbb{Z}\Gamma(G/H) \cap \Lambda^+. \quad (2.6)$$

Therefore, if Γ is saturated, the G -algebra $\mathbb{C}[G/\widehat{H}]$ is isomorphic to the graduate algebra associated to the maximal filtration $\mathfrak{F}_\mu$. Hence $\text{Spec } \mathbb{C}[G/\widehat{H}]$ is a deformation of $\text{Spec } \mathbb{C}[G/H]$.

This can be applied in our setting. Indeed, if Γ is free and saturated and X_0 is the affine multicone defined above, then a spherical G -subvariety X_1 of V is a deformation of X_0 if and only if its open G -orbit is a deformation of $G.v_\lambda$ (see [BK]).

3 The tangent space of the invariant Hilbert scheme

The purpose of this section is the proof of Theorem 2.2. We first compute the T_{ad} -weights of the tangent space of $\text{Hilb}_\Gamma^G(V)$ at the multicone X_0 $T_{X_0}\text{Hilb}_\Gamma^G(V)$ and then prove its multiplicity freeness.

Recall that $\lambda_1, \dots, \lambda_s$ are linearly independent dominant weights which generate a saturated submonoid Γ and that V is the G -module $V(\lambda_1) \oplus \dots \oplus V(\lambda_s)$.

Since $\text{codim}_{X_0}(X_0 \setminus G.v_\lambda) \geq 2$ (see (2.5)) we can use the following characterisation of the tangent space of the invariant Hilbert scheme at the multicone X_0 given by Proposition 1.15 of [AB]

$$T_{X_0}\text{Hilb}_\Gamma^G(V) \cong (V/\mathfrak{g}.v_\lambda)^{G_{v_\lambda}} \quad (3.1)$$

where $\mathfrak{g}$ denotes the Lie algebra of G and $v_{\underline{\lambda}} = v_{\lambda_1} + \dots + v_{\lambda_s}$, with v_{λ_i} a highest weight vector in $V(\lambda_i)$.

3.1 Action of the adjoint torus - its weights

Consider the following action of the adjoint torus $T_{\text{ad}} = T/Z(G)$ on V defined on every T -weight space $V(\lambda_i)_{\mu}$ by

$$t.v = \lambda_i(t)\mu(t)^{-1}v \quad (3.2)$$

(see Section 2.1 in [AB]). It yields a T_{ad} -action on the vector space $(V/\mathfrak{g}.v_{\underline{\lambda}})^{G_{v_{\underline{\lambda}}}}$. This action is often called the *normalised action* of T_{ad} on $(V/\mathfrak{g}.v_{\underline{\lambda}})^{G_{v_{\underline{\lambda}}}}$. It corresponds to the differential of the T_{ad} -action on the invariant Hilbert scheme $\text{Hilb}_{\Gamma}^G(V)$ defined in (1.3) (see *loc. cit.*).

The stabiliser $G_{v_{\underline{\lambda}}}$ is generated by its maximal torus, intersection of the kernels of the characters λ_i , and its unipotent part. The T_{ad} -module $(V/\mathfrak{g}.v_{\underline{\lambda}})^{G_{v_{\underline{\lambda}}}}$ is thus equal to

$$(V/\mathfrak{g}.v_{\underline{\lambda}})_{\langle \lambda_1, \dots, \lambda_s \rangle}^{\mathfrak{g}_{v_{\underline{\lambda}}}}, \quad (3.3)$$

that is, the subspace spanned by the T_{ad} -semi-invariant classes in $(V/\mathfrak{g}.v_{\underline{\lambda}})^{\mathfrak{g}_{v_{\underline{\lambda}}}}$ whose T_{ad} -weights are integral linear combinations of the weights λ_i .

Note that representatives of T_{ad} -weight vectors of $(V/\mathfrak{g}.v_{\underline{\lambda}})^{\mathfrak{g}_{v_{\underline{\lambda}}}}$ with weight γ can be chosen in $\oplus_i V(\lambda_i)_{\lambda_i - \gamma}$.

Theorem 3.1. *Let $\lambda_1, \dots, \lambda_s$ be linearly independent dominant weights generating a saturated cone Γ . Consider the G -module $V = \oplus_1^s V(\lambda_i)$. The T_{ad} -weights of $(V/\mathfrak{g}.v_{\underline{\lambda}})^{G_{v_{\underline{\lambda}}}}$ for the normalised action belong to Table 1.*

How to read Table 1. A T_{ad} -weight γ is an element of the root lattice. In the table it is written as linear combination of simple roots. The simple roots are labelled according to the type of the support of the T_{ad} -weight γ .

Corollary 3.2. *The T_{ad} -weights of $(V/\mathfrak{g}.v_{\underline{\lambda}})^{G_{v_{\underline{\lambda}}}}$ are spherical roots.*

Proof. From the list stated in Table 1, one checks easily that the elements of Σ are indeed spherical roots (compare with Table 1 in [W] which lists all the spherical roots). $\square$

The following is devoted to the proof of Theorem 3.1.

Lemma 3.3. *Let $[v] \neq 0$ be a T_{ad} -weight vector in $(V/\mathfrak{g}.v_{\underline{\lambda}})^{G_{v_{\underline{\lambda}}}}$. Then there exists a pair (α, β) of positive roots with α simple such that (up to a rescaling of v)*

$$X_{\alpha}.v \neq 0 \quad \text{and} \quad X_{\alpha}.v = X_{-\beta}.v_{\underline{\lambda}}. \quad (3.4)$$

Table 1: T_{ad} -weights in $(V/\mathfrak{g}.v_{\underline{\lambda}})^{G_{v_{\underline{\lambda}}}}$.

Type of support	Weight
$A_1 \times A_1$	$\alpha + \alpha'$
A_n	$\alpha_1 + \dots + \alpha_n, n \geq 2$ $2\alpha, n = 1$ $\alpha_1 + 2\alpha_2 + \alpha_3, n = 3$
$B_n, n \geq 2$	$\alpha_1 + \dots + \alpha_n$ $2\alpha_1 + \dots + 2\alpha_n$ $\alpha_1 + 2\alpha_2 + 3\alpha_3, n = 3$
$C_n, n \geq 3$	$\alpha_1 + 2\alpha_2 + \dots + 2\alpha_{n-1} + \alpha_n$
$D_n, n \geq 4$	$2\alpha_1 + \dots + 2\alpha_{n-2} + \alpha_{n-1} + \alpha_n$ $\alpha_1 + 2\alpha_2 + 2\alpha_3 + \alpha_4, n = 4$ $\alpha_1 + 2\alpha_2 + \alpha_3 + 2\alpha_4, n = 4$
F_4	$\alpha_1 + 2\alpha_2 + 3\alpha_3 + 2\alpha_4$
G_2	$4\alpha_1 + 2\alpha_2$ $\alpha_1 + \alpha_2$

Proof. The class $[v]$ being different from 0, the vector v is not a linear combination of highest weight vectors v_{λ_i} . Hence, there exists $\alpha \in S$ such that $X_{\alpha}.v \neq 0$. Since $\mathfrak{g}_{v_{\underline{\lambda}}}.v \subset \mathfrak{g}.v_{\underline{\lambda}}$, we have $X_{\alpha}.v \in \mathfrak{g}.v_{\underline{\lambda}}$.

Assume $X_{\alpha}.v$ is a linear combination of highest weight vectors. The simple root α is thus the T_{ad} -weight of $[v]$ and v can be chosen in $\oplus_i V(\lambda_i)_{\lambda_i - \alpha}$. Hence, there exists j such that $(\lambda_j, \alpha) \neq 0$. Because of saturation, by Lemma 2.1, such a weight λ_j is unique, since α is an integral combination of the weights λ_i . This means that the i -th component of v in $V(\lambda_i)$ is zero, for all $i \neq j$. Hence, $v \in \mathbb{C}X_{-\alpha}.v_{\underline{\lambda}}$ which is not possible because $[v] \neq 0$. Therefore, there exists $\beta \in \Phi^+$ such that $X_{\alpha}.v \in \mathbb{C}X_{-\beta}.v_{\underline{\lambda}}$. $\square$

Remark 3.1. The sum $\alpha + \beta$ of any pair (α, β) of positive roots (with α not necessarily simple) which satisfies condition (3.4) is equal to the weight of $[v]$.

Proposition 3.4. *Let γ be a T_{ad} -weight of some $(V/\mathfrak{g}.v_{\underline{\lambda}})^{G_{v_{\underline{\lambda}}}}$. Suppose there exists a simple root δ in the support of γ such that $\gamma - \delta \notin \Phi$. Then $(\gamma, \delta) \geq 0$. Moreover if $(\gamma, \delta) = 0$ then $(\lambda_i, \delta) = 0$ for all i .*

Proof. Let us prove the two assertions together.

We shall proceed by contradiction. Suppose the root δ is such that $(\gamma, \delta) \leq 0$ and there exists j such that $(\lambda_j, \delta) \neq 0$ (that is obviously true

if $(\gamma, \delta) \neq 0$.

By Lemma 2.1 there exist a simple root α and a weight λ_k with $(\gamma, \alpha) > 0$, $(\lambda_k, \alpha) > 0$ and $(\lambda_i, \alpha) = 0$ for all $i \neq k$. Since λ_k occurs with a positive coefficient in the writing of γ as an integral linear combination of the weights λ_i , if (λ_k, δ) is nonzero (λ_i, δ) can not be zero for all $i \neq k$. Hence, we can suppose $(\lambda_j, \alpha) = 0$.

Let $[v] \in (V/\mathfrak{g}.v_\lambda)^{G_{v_\lambda}}$ be nonzero of T_{ad} -weight γ . Consider $v_j \in V(\lambda_j)$ the j -th component of some representative v of $[v]$. Note that we have $(\lambda_j - \gamma, \alpha) < 0$. Hence, if $v_j \neq 0$ then $X_\alpha.v_j \neq 0$, identity (3.4) holds with the pair $(\alpha, \gamma - \alpha)$ and in particular $X_\alpha.v_j = X_{-\gamma+\alpha}.v_{\lambda_j}$. This implies that v_j is proportional to $X_{-\alpha}X_{-\gamma+\alpha}.v_{\lambda_j}$ which equals $X_{-\gamma}.v_{\lambda_j}$ in case $\gamma \in \Phi$ and 0 otherwise. In any case, we thus have: $v_j \in \mathfrak{g}.v_{\lambda_j}$ hence we can choose $v' \in [v]$ such that $v'_j = 0$.

Take α' and β' that fulfill (3.4) for the vector v' , we have: $(\lambda_j, \gamma - \alpha') = 0$. Since $\delta \in \text{Supp } \gamma$ and $(\lambda_j, \delta) \neq 0$, the only possibility is that $\delta = \alpha'$. But $\gamma - \delta$ is not a root, hence (3.4) can not hold with δ . This yields the contradiction. $\square$

Lemma 3.5. *Let γ be a T_{ad} -weight of $(V/\mathfrak{g}.v_\lambda)^{G_{v_\lambda}}$ and L be the standard Levi subgroup L associated to the set of simple roots $\text{Supp } \gamma$. Denote W the L -submodule of V generated by $v_{\pi(\lambda)}$. Then*

$$(V/\mathfrak{g}.v_\lambda)_\gamma^{G_{v_\lambda}} \cong (W/\mathfrak{l}.v_{\pi(\lambda)})_\gamma^{L_{v_{\pi(\lambda)}}}. \quad (3.5)$$

as T_{ad} -modules, where π is the projection of Λ onto the weight lattice of L .

Proof. The isomorphism of the lemma is given by sending $[v] \in V/\mathfrak{g}.v_\lambda$ to the class of v in $W/\mathfrak{l}.v_\lambda$ where $v \in \oplus_i V(\lambda_i)_{\lambda_i - \gamma}$. $\square$

Lemma 3.6. *Let γ be a T_{ad} -weight of some $(V/\mathfrak{g}.v_\lambda)^{G_{v_\lambda}}$. Suppose γ is not a root. Consider α and β two positive roots which fulfill condition (3.4). If $\beta = \beta_1 + \beta_2$ with β_1 and β_2 positive roots then $\alpha + \beta_1$ or $\alpha + \beta_2$ is a root.*

Proof. Let $v \in V$ satisfy (3.4) with the given roots α and β , in particular $\gamma = \alpha + \beta$. Assume $\alpha + \beta - \beta_i$ is not a root for $i = 1, 2$. Since $X_{\beta_i}.v \in \mathfrak{g}.v_\lambda$ we have: $X_{\beta_i}.v = 0$ for $i = 1, 2$ hence $X_\beta.v = 0$. It follows along with $\gamma \notin \Phi$ and the property (3.4): $0 = X_\alpha X_\beta.v = X_\beta X_\alpha.v = X_\beta X_{-\beta}.v_\lambda$. In particular, we have: $X_{-\beta}.v_\lambda = 0$ which contradicts $0 \neq X_\alpha.v = X_{-\beta}.v_\lambda$ given by (3.4). $\square$

Lemma 3.7. *Let $[v]$ be a nonzero semi-invariant class in $(V/\mathfrak{g}.v_\lambda)_{v_\lambda}^G$ and (α, β) pair of positive roots, with α simple, fulfilling condition (3.4) for the vector v . If α and β are orthogonal and $\alpha + \beta$ is not a root, namely the pair (α, β) generates a root subsystem of type $A_1 \times A_1$, then there exists a unique j such that λ_j is not orthogonal to $\alpha + \beta$.*

Proof. Consider the root subsystem Φ' generated by α and β and apply Lemma 3.5. $\square$

Lemma 3.8. *Let γ be the weight of a nonzero class $[v] \in (V/\mathfrak{g}.v_\lambda)^G_{v_\lambda}$. If one of the following conditions is fulfilled, there exists a unique j such that λ_j is not orthogonal to γ .*

- (i) *There exists a unique root $\delta_0 \in \text{Supp } \gamma$ such that $(\gamma, \delta_0) \neq 0$ and there exist no root $\delta \in \text{Supp } \gamma$ such that $(\gamma, \delta) = 0$ and $\gamma - \delta \in \Phi$.*
- (ii) *There exists a unique simple root α such that $(\gamma, \alpha) > 0$ and $\gamma - \alpha \in \Phi$. Moreover, for all $\delta \in \text{Supp } \gamma \setminus \{\alpha\}$ we have: $(\gamma, \delta) \leq 0$ and $\gamma - \delta \notin \Phi$.*

Proof. By saturation and Proposition 3.4. $\square$

Proof of Theorem 3.1. Pick a T -weight γ of $(V/\mathfrak{g}.v_\lambda)^{G_{v_\lambda}}$. We know by Lemma 3.3 that the weight γ is a sum of two positive roots, say α and β with α simple. If α is orthogonal to the support of β , then β is also simple by Lemma 3.6.

This allows to reduce to the case of a simple group G .

Finally most of the choices for $\gamma = \alpha + \beta$ can be ruled out: in some cases we can use Lemma 3.6, in other cases it is possible to prove that there is only one dominant weight λ_i not orthogonal to γ (applying Lemma 3.7 and Lemma 3.8) hence we can use Lemma 3.5 and Theorem 1.1 in [J].

We end up with the list of Table 1.

3.2 Multiplicity freeness

Theorem 3.9. *Each T_{ad} -weight space of $(V/\mathfrak{g}.v_\lambda)^{G_{v_\lambda}}$ is one dimensional.*

We first work out the case of irreducible modules V then in combination with Lemma 3.5 we get to the general case.

Proposition 3.10. *Suppose V is an irreducible G -module, namely $V = V(\lambda)$ for a dominant weight λ . Then the T_{ad} -weight spaces of $(V/\mathfrak{g}.v_\lambda)^{G_{v_\lambda}}$ are one dimensional.*

The previous statement was already proved in [J]. We present here a more concise proof.

Proof. From Table 1, we know that if the weight γ is not a root then there exists exactly one simple root δ such that $(\gamma, \delta) \neq 0$. Hence, by Proposition 3.4, the dominant weight λ is (up to a scalar) the fundamental weight associated to δ . Moreover, for such a root δ , we can check that the coefficient

corresponding to δ in the writing of γ is 2 or 3. Therefore, if this coefficient is 2, the following vectors are clearly basis vectors of the weight space $V_{\gamma-\lambda}$:

$$X_{-\gamma+\nu}X_{-\nu}.v_\lambda,$$

where $\{\nu, \gamma - \nu\}$ runs over the 2-set of positive roots such that ν and $\gamma - \nu$ contain δ in their support. This holds also in case the coefficient of δ is 3. Indeed, we have then $\gamma = \alpha_1 + 2\alpha_2 + 3\alpha_3$ of type B_3 . Consider the vectors $X_{-\gamma+\nu_1+\nu_2}X_{-\nu_1}X_{-\nu_2}.v_\lambda$ with $\{\nu_1, \nu_2, \gamma - (\nu_1 + \nu_2)\}$ running over 3-sets of positive roots ν_1, ν_2 and $\gamma - (\nu_1 + \nu_2)$ which all contain δ in their support. Then it is easy to see that these vectors are linear combinations of the above vectors $X_{-\gamma+\nu}X_{-\nu}.v_\lambda$.

Applying now the equalities $X_\nu.v = 0$ for all $\nu \in \text{Supp } \gamma \setminus \{\delta\}$, we obtain the proposition.

If γ is a root then we are left with $\gamma = \alpha_1 + \dots + \alpha_n$, $\gamma = \alpha_1 + 2(\alpha_2 + \dots) + \alpha_n$ or $\gamma = \alpha_1 + 2\alpha_2 + 3\alpha_3 + 2\alpha_4$ respectively of type A_n , C_n and F_4 . Indeed, type B_n (resp. G_2) is easily ruled out since the weight space $V(\lambda)_{\lambda-\gamma}$ is spanned by $X_{-\gamma}.v_\lambda$ (resp. $\gamma = \omega_2 - \omega_1$ is not up to a scalar a dominant weight). For the possible remaining weights γ , we take as basis vectors of $V(\lambda)_{\lambda-\gamma}$ the same vectors $X_{-\gamma+\nu}X_{-\nu}.v_\lambda$ as above ($\delta = \alpha_1, \alpha_2$ or α_4 respectively) along with $X_{-\gamma}.v_\lambda$ and conclude like before. $\square$

Proof. Let γ be a T_{ad} -weight of $(V/\mathfrak{g}.v_\lambda)^{G_{v_\lambda}}$.

We first consider the case where G is of type G_2 and $\gamma = \alpha_1 + \alpha_2$, we find as single possibility for $\underline{\lambda}$: $\lambda_1 = \omega_1$ and $\lambda_2 = \omega_2$ and the tangent space is one dimensional.

Suppose now that if $\gamma = \alpha_1 + \alpha_2$ then its support is not of type G_2 . From Table 1, we get:

$$(\gamma, \alpha) \geq 0 \text{ for every } \alpha \in \text{Supp } \gamma.$$

Note that this can be proved also by means of Proposition 3.4.

One can check in Table 1 that there are either one or two simple roots δ such that $(\gamma, \delta) > 0$. Such roots are necessarily in $\text{Supp } \gamma$.

Consider first the case where there exists a unique simple root, say δ , such that $(\gamma, \delta) > 0$. Hence, because of saturation, there exists a unique dominant weight λ_i such that $(\lambda_i, \delta) \neq 0$. By assumption we have: $(\gamma, \alpha) = 0$ for every $\alpha \in \text{Supp } \gamma \setminus \{\delta\}$.

If $(\underline{\lambda}, \alpha) = 0$ for every simple root $\alpha \in \text{Supp } \gamma \setminus \{\delta\}$, we have:

$$\oplus_j V(\lambda_j)_{\lambda_j-\gamma} = V(\lambda_i)_{\lambda_i-\gamma}.$$

and the theorem follows from the reduction to the Levi subgroup associated to the support of γ and Proposition 3.10.

Suppose that $(\underline{\lambda}, \alpha) \neq 0$ for a simple root $\alpha \in \text{Supp } \gamma \setminus \{\delta\}$. By Proposition 3.4, the root α is such that $\gamma - \alpha \in \Phi$. Note that the weight γ is obviously a root and the root α is unique. By Table 1, we have even more precisely that $\gamma = \alpha_1 + \dots + \alpha_n$, $\gamma = \alpha_1 + 2(\alpha_2 + \dots) + \alpha_n$ or $\gamma = \alpha_1 + 2\alpha_2 + 3\alpha_3 + 2\alpha_4$ respectively of type B_n , C_n and F_4 . The root α is respectively α_n , α_1 and α_3 . Then $\underline{\lambda}$ have one of the following shapes (after restriction to the Levi subgroup associated to the support of γ):

$$(\omega_\delta, a_1\omega_\alpha, \dots, a_r\omega_\alpha) \quad \text{or} \quad (\omega_\delta + b\omega_\alpha, b_1\omega_\alpha, \dots, b_r\omega_\alpha)$$

where the a_i 's, b and the b_j 's are positive integers and $r \geq 1$.

Note that the case of type F_4 is special. Only the first shape of $\underline{\lambda}$ is possible a priori, with $r = 1$, but in this case also the tangent space is 0-dimensional. We postpone it in a separated lemma below.

In type B_n and C_n the weight spaces $V(\lambda_i)_{\lambda_i - \gamma}$ are clearly spanned by $X_{-\gamma} \cdot v_{\lambda_i}$ for any positive integer a_i, b_i ($i = 1, \dots, r$). Let $[v]$ be a T_{ad} -vector of weight γ with $v \in \oplus_i V(\lambda_i)_{\lambda_i - \gamma}$. We can thus suppose all its components but the first one to be 0. Indeed, one can take as new representative for $[v]$, the vector $v - kX_{-\gamma+\delta} \cdot v_{\underline{\lambda}}$ if one of the v_j ($j > 1$) is such that $0 \neq v_j = kX_{-\gamma+\delta} \cdot v_{\lambda_j}$ (k being a non-zero scalar).

If $\lambda_1 = \omega_\delta$ with $\gamma = \alpha_1 + \dots + \alpha_n$ (here $\delta = \alpha_1$) then the tangent space is clearly one dimensional and we can take as T_{ad} -weight vector the class of $(0, \dots, 0, X_{-\gamma} \cdot v_{\underline{\lambda}})$. Remark that only in this case the weight space $V(\lambda_1)_{\lambda_1 - \gamma}$ is one dimensional. For the other cases under consideration, one basis of $V(\lambda_1)_{\lambda_1 - \gamma}$ is given by the following vectors:

$$X_{-\gamma} \cdot v_{\lambda_1}, X_{-\gamma+\nu} X_{-\nu} \cdot v_{\lambda_1}$$

with $\{\delta\} \subset \text{Supp } \nu \subset \text{Supp } \gamma \setminus \{\alpha\}$.

Recall that $X_\nu \cdot v = 0$ for all $\nu \in \text{Supp } \gamma \setminus \{\alpha, \delta\}$ since $\gamma - \nu \notin \Phi$. Writing v as a linear combination of the above basis vectors and using these equalities, we obtain that the tangent space is multiplicity free.

Consider now the case where there exist two distinct simple roots, say δ_1 and δ_2 , such that $(\gamma, \delta_1) > 0$ and $(\gamma, \delta_2) > 0$. According to Table 1, this assumption is only valid for $\gamma = \alpha_1 + \dots + \alpha_n$ of type A_n . After reduction to the Levi subgroup associated to $\text{Supp } \gamma$, we get as possibilities for $\underline{\lambda}$:

$$\omega_1 + \omega_n, (\omega_1, \omega_n), (a\omega_1 + \omega_n, b\omega_1), (a\omega_1 + \omega_n, b\omega_1, c\omega_1)$$

where a, b, c are positive integers. Note that the last case can only occur in type D and E. In all these cases, one shows similarly as before that the components in $V(\lambda_i)$, $i = 2, 3$, of a T_{ad} -weight vector can be taken to be equal to 0. Then straightforward computations lead to the proposition. $\square$

Lemma 3.11. *If G is of type F_4 , $\lambda_1 = \omega_4$ and $\lambda_2 = a\omega_3$ with $a > 0$, then the space $(V/\mathfrak{g}.v_\lambda)^{G_{v_\lambda}}$ is trivial.*

Proof. We can take $v_2 = 0$ since it should be in $(V(\lambda_2)/\mathfrak{g}.v_{\lambda_2})^{G_{v_\lambda}}$ and in fact in $(V(\lambda_2)/\mathfrak{g}.v_{\lambda_2})^{G_{v_{\lambda_2}}}$ which is trivial as proved previously. We have also $v_1 \in (V(\lambda_1)/\mathfrak{g}.v_{\lambda_1})^{G_{v_{\lambda_1}}}$ and recall that in this case the space is one dimensional. Suppose $v_1 \neq 0$ then since $X_{\alpha_3}.v_1 \neq 0$, it should be (up to a scalar) equal to $X_{-\gamma+\alpha_3}.v_{\lambda_1}$. Hence $X_{\alpha_3}.v_2$ (which is 0) is not equal to $X_{-\gamma+\alpha_3}.v_{\lambda_2}$. $\square$

4 Spherical systems

After having recalled the axiomatic definition of a spherical system given in [L01], we show how we can associate naturally to the given monoid Γ a spherical system. Further, we prove in the last paragraph that Γ fulfills a certain regularity property.

4.1 Definition of spherical systems

Definition 4.1. A *spherical system* for G is a triple consisting of a subset S^p of simple roots, a set Σ of spherical roots for G (namely T -weights that are the spherical root of a rank one wonderful G -variety) and a set $\mathbf{A}$ endowed with a map $\rho: \mathbf{A} \rightarrow \Xi^*$, where $\Xi = \mathbb{Z}\Sigma$, which satisfies the following properties.

(A1) For every $D \in \mathbf{A}$ and $\gamma \in \Sigma$ we have $\langle \rho(D), \gamma \rangle \leq 1$, and if $\langle \rho(D), \gamma \rangle = 1$ then $\gamma \in S \cap \Sigma$.

(A2) For every $\alpha \in \Sigma \cap S$, $\mathbf{A}$ contains two elements, D_α^+ and D_α^- , such that $\langle \rho(D_\alpha^\pm), \alpha \rangle = 1$. Moreover $\langle \rho(D_\alpha^+), \gamma \rangle + \langle \rho(D_\alpha^-), \gamma \rangle = \langle \gamma, \alpha^\vee \rangle$, for every $\gamma \in \Sigma$.

(A3) The set $\mathbf{A}$ is the union of $\{D_\alpha^+, D_\alpha^-\}$ for all $\alpha \in \Sigma \cap S$.

(Σ1) If $2\alpha \in \Sigma \cap 2S$ then $\frac{1}{2}\langle \alpha^\vee, \gamma \rangle$ is a nonpositive integer for every $\gamma \in \Sigma \setminus \{2\alpha\}$.

(Σ2) If $\alpha, \beta \in S$ are orthogonal and $\alpha + \beta \in \Sigma$ or $\frac{1}{2}(\alpha + \beta) \in \Sigma$ then $(\gamma, \alpha^\vee) = (\gamma, \beta^\vee)$ for every $\gamma \in \Sigma$.

(S) For every $\gamma \in \Sigma$, there exists a rank one wonderful G -variety X with γ as spherical root and S^p equal to the set of simple roots associated to P_X .

Remark 4.1. The last axiom (S) of spherical system is equivalent, in type different from F , to the following. For all $\gamma \in \Sigma$,

$$\{\alpha \in S : (\gamma, \alpha) = 0 \text{ and } \gamma - \alpha \notin \Phi\} \cap \text{Supp } \gamma \subseteq S^p \subseteq \{\alpha \in S : (\gamma, \alpha) = 0\}. \quad (4.1)$$

4.2 The spherical system attached to Γ

Let Γ be a saturated cone generated by linearly independent dominant weights $\lambda_1, \dots, \lambda_s$. Let V be the finite dimensional G -module $\oplus_i V(\lambda_i)$. We denote by S^p be the subset of simple roots orthogonal to λ_i for all i and by Σ be the set of T_{ad} -weights in $(V/\mathfrak{g}.v_\lambda)^{G_{v_\lambda}}$.

Theorem 4.1. *The triple $(S^p, \Sigma, \emptyset)$ is a spherical system for G .*

The two following results are directly involved in the proof of Theorem 4.1 and Proposition 4.4. The second one is a consequence of the first one and of Theorem 2.2.

Lemma 4.2. *Let $[v] \in (V/\mathfrak{g}.v_\lambda)^{G_{v_\lambda}}$ be of weight γ .*

- (1) *If $\gamma = 2\alpha$ with $\alpha \in S$, there exists a unique j such that $(\lambda_j, \alpha) \neq 0$. Moreover, $(\lambda_j, \alpha^\vee)$ is even.*
- (2) *If $\gamma = \alpha + \beta$ with $\alpha, \beta \in S$ and $(\alpha, \beta) = 0$, there exists a unique j such that $(\lambda_j, \alpha).(\lambda_j, \beta) \neq 0$. Moreover, $(\lambda_j, \alpha^\vee) = (\lambda_j, \beta^\vee)$ and $(\lambda_i, \alpha) = (\lambda_i, \beta) = 0$ for all $i \neq j$.*

Proof. (1) Since $v \in \oplus_i V(\lambda_i)_{\lambda_i - 2\alpha}$, one of the dominant weights λ_i , say λ_j , is such that $(\lambda_j, \alpha^\vee) \geq 2$. Recall that the weights of $(V/\mathfrak{g}.v_\lambda)^{G_{v_\lambda}}$ are integral linear combinations of the weights λ_i . Since the weight of $[v]$ is equal up to a scalar to a simple root, the coefficients involved in its writing as linear combination of the weights λ_i are all negative except that of λ_j . Because of saturation, we thus get: $(\lambda_i, \alpha) \neq 0$ if and only if $i = j$. Moreover, we have that $(\lambda_j, \alpha^\vee)$ divides $(2\alpha, \alpha^\vee) = 4$. With the above inequality this implies that $(\lambda_j, \alpha^\vee)$ is even.

(2) The given simple roots α, β generate a root system of type $A_1 \times A_1$. Since $v \in \oplus_i V(\lambda_i)_{\lambda_i - \gamma}$, there exists λ_j such that $(\lambda_j, \alpha).(\lambda_j, \beta) \neq 0$. By saturation, λ_j is the unique dominant weight which satisfies this inequality. Therefore, the coordinates $v_i \in V(\lambda_i)$ of v are all 0 except v_j . The only two pairs which can fulfill condition ((3.4)) are (α, β) and (β, α) . Suppose $0 \neq X_\alpha.v = X_{-\beta}.v_\lambda$, we then have: $0 = X_\alpha.v_i = X_{-\beta}.v_{\lambda_i}$ for $i \neq j$. From $0 \neq (\lambda_j, \beta^\vee)v_{\lambda_j} = X_\beta X_\alpha.v_j = X_\alpha X_\beta.v_j = (\lambda_j, \alpha^\vee)v_{\lambda_j}$, we get also that $X_\beta.v_j \neq 0$, hence $(\lambda_i, \alpha) = (\lambda_i, \beta) = 0$ for all $i \neq j$ and $(\lambda_j, \alpha^\vee) = (\lambda_j, \beta^\vee)$. $\square$

Corollary 4.3. *The set Σ of T_{ad} -weights of $(V/\mathfrak{g}.v_\lambda)^{G_{v_\lambda}}$ satisfies the following properties.*

- ($\Sigma 1$) *Let α be a simple root such that $2\alpha \in \Sigma$. Then $(\gamma, \alpha^\vee) \in 2\mathbb{Z}_{\leq 0}$ for all $\gamma \in \Sigma \setminus \{2\alpha\}$.*
- ($\Sigma 2$) *Let α and β be orthogonal simple roots. If $\alpha + \beta \in \Sigma$ then $(\gamma, \alpha^\vee) = (\gamma, \beta^\vee)$ for every $\gamma \in \Sigma$.*

Proof. Suppose there exists a weight $\gamma \in \Sigma \setminus \{2\alpha\}$ such that $(\gamma, \alpha) > 0$. By the above argument, we have: $(\gamma, \alpha^\vee) \geq 2$. Hence in case γ is a root, so are $\gamma - \alpha$ and $\gamma - 2\alpha$, but according to the list given in Theorem 2.2, this can not be. Consider now, the case where γ is not a root. Checking again the list of possible T_{ad} -weights, one obtains as single possibility $\gamma = \alpha_{i-1} + 2\alpha_i + \alpha_{i+1}$ with $\alpha_i = \alpha$. Hence we have in this case a dominant weight which is not orthogonal to α_{i-1} . Applying Proposition 3.4 with the simple root α_{i-1} , this yields a contradiction. The first assertion follows.

The second assertion is contained in Lemma 4.2(2). $\square$

Proof of Theorem 4.1. Since by Lemma 3.3 the set Σ does not contain any simple root we need to prove only axioms $(\Sigma 1)$, $(\Sigma 2)$ and (S) . The first two follow from Corollary 4.3 and the latter from Proposition 3.4 and Remark 4.1. The axiom (S) for the spherical root with support of type F follows from Lemma 3.11.

4.3 Regularity of Γ

Let us first introduce the the set of *colours* Δ associated to a spherical system $(S^p, \Sigma, \mathbf{A})$ following [L01].

The set Δ is the disjoint union

$$\Delta = \mathbf{A} \cup \Delta^{a'} \cup \Delta^b \quad (4.2)$$

where $\Delta^{a'}$ (resp. Δ^b) contains an element D_α for each $\alpha \in S$ such that $2\alpha \in \Sigma$ (resp. $\alpha \notin S^p \cup \Sigma \cup \frac{1}{2}\Sigma$). If $\alpha + \alpha' \in \Sigma$ or $\frac{1}{2}(\alpha + \alpha') \in \Sigma$ then we put $D_\alpha = D_{\alpha'}$. Note that Δ is indeed a disjoint union because of the conditions (Σ_1) and (Σ_2) of spherical systems.

Denote ω_α the fundamental weight associated to a simple root α . Then one defines a map $\sigma : \mathbb{Z}\Delta \rightarrow \Lambda$ as follows (see [F])

$$D \mapsto \begin{cases} \sum \omega_\alpha & \text{if } D \in \mathbf{A} \cup \Delta^b \text{ and } D = D_\alpha \\ 2\omega_\alpha & \text{if } D \in \Delta^{a'} \text{ and } D = D_\alpha \end{cases} . \quad (4.3)$$

Proposition 4.4. *Consider the spherical system associated to a free and saturated monoid $\Gamma = \langle \lambda_1, \dots, \lambda_s \rangle$. Let Δ be its set of colours.*

(i) *We can regard the cone $\mathbb{N}\Delta$ generated by the set of colours as a subset of Λ^+ . The cone Γ is then included in the cone $\mathbb{N}\Delta$.*

(ii) *Each colour of Δ occurs in the linear combination of at least one weight λ_i .*

The second assertion of the proposition can be called *property of regularity*.

Proof. Since the set $\mathbf{A}$ of the spherical system is empty, the map σ defined above is injective.

The image of a colour in Λ^+ is equal to $2\omega_\alpha$ for all simple roots α with $2\alpha \in \Sigma$, $\omega_\alpha + \omega_\beta$ for all orthogonal simple roots α, β with $\alpha + \beta \in \Sigma$ and ω_α for the remaining simple roots $\alpha \notin S^p$. Since $S^p = \{\alpha \in S : (\lambda_i, \alpha) = 0 \forall i\}$ the assertion follows from Lemma 4.2. $\square$

5 The affine multicone over a wonderful variety

5.1 The spherical system of a wonderful variety

Let X be a wonderful G -variety. The set Σ_X of its spherical roots is a basis of the lattice Ξ_X of weights of B -semi-invariant rational functions on X .

A colour of X is a B -stable not G -stable prime divisor in X , namely the closure of a colour (a B -stable prime divisor) of the open G -orbit. Let Δ_X denote the set of colours of X .

If P_X denotes the standard parabolic subgroup equal to the stabiliser of the colours of X , the closed orbit of X is isomorphic to G/P_X . Let S_X^p denote the set of simple roots associated to P_X .

Let $\mathbf{A}_X$ denote the set of colours D such that there is a simple root α in Σ_X with D not P_α -stable.

Define the map $\rho_X : \Delta_X \rightarrow \Xi_X^*$ by $\langle \rho_X(D), \gamma \rangle = v_D(f_\gamma)$, where v_D is the valuation associated to the divisor D and f_γ is the B -semi-invariant rational function of weight γ (uniquely determined up to a scalar).

Then the triple $S_X^p, \Sigma_X, \mathbf{A}_X$ is a spherical system for G and Δ_X is the set of colours associated to it ([L01] Section 7).

5.2 The open orbit

Recall the notation of Paragraph 4.2. Let $(S^p, \Sigma, \emptyset)$ be the spherical system for G attached to Γ .

We shall assume in the following that there exists an associated wonderful subgroup N of G , that is the generic stabiliser of a wonderful G -variety X with given spherical system $(S^p, \Sigma, \emptyset)$.

In Section 6 we will prove the existence of such a wonderful subgroup for all spherical systems arising from a free and saturated cone.

The main purpose of this section is to prove the second assertion of Theorem 2.3 under this existence assumption, that is

Proposition 5.1. *There exists a subgroup H of N , spherical in G , such that the homogeneous space G/H is quasi-affine with weight monoid Γ . We may regard G/H in V then its closure X_1 in V is a deformation of X_0 .*

We need a reformulation of Proposition 4.4 in terms of regular functions on G . Therefore, we shall recall an equivalent definition of the map σ given in (4.3). We consider here the opposite Borel subgroup B^- of B . We can suppose that the set $B^- N$ is open in G .

The inverse image of a B^- -stable prime divisor of G/N through the quotient map $G \rightarrow G/N$ admits an equation in $\mathbb{C}[G]$ which is B^- -left-semi-invariant and N -right-semi-invariant or for short a $B^- \times N$ -eigenvector of $\mathbb{C}[G]$. This defines an injective map on the set of colours of G/N

$$\sigma' : \mathbb{Z}\Delta \hookrightarrow \Lambda \times \Xi(N) \quad (5.1)$$

where $\Xi(N)$ is the lattice of N -weights.

Consider now the composition of the map σ' of (5.1) with the projection onto Λ . Such a map

$$\sigma'' : \mathbb{Z}\Delta \rightarrow \Lambda \quad (5.2)$$

is indeed the map $-\sigma$ (see [F]).

We thus get the reformulation of Proposition 4.4.

Proposition 5.2. (i) *For any weight χ in Γ , there exists an unique $B^- \times N$ -eigenvector of $\mathbb{C}[G]$ of B^- -weight $-\chi$.*

(ii) *Let f_i be the $B^- \times N$ -eigenvector of $\mathbb{C}[G]$ of weight $-\lambda_i$ for $i = 1, \dots, s$. Then the function $\prod_{j=1, \dots, s} f_j$ vanishes on $G \setminus B^- N$.*

Proof of Proposition 5.1 Via the isomorphism of G -modules $\mathbb{C}[G] \cong \bigoplus_{\lambda \in \Lambda^+} V(\lambda)^* \otimes V(\lambda)$ the function f_i corresponds to $\eta_i \otimes v_i \in V(\lambda_i)^* \otimes V(\lambda_i)$, where η_i is a B^- -eigenvector and v_i a N -eigenvector.

Consider the point $x = ([v_1], \dots, [v_s])$ in the product of the projective spaces $\mathbb{P}(V(\lambda_i))$. It is therefore N -stable. Further, from the second assertion of Proposition 5.2, we have: $N \subset G_x \subset N_G(N)$.

We can suppose $N = N_G(N)$. Indeed, the normaliser $N_G(N)$ acts on the set of colours of G/N preserving the associated functionals on the lattice Ξ generated by Σ . Since in our case there are no spherical roots equal to simple roots, there can not be distinct colours with the same functional, hence the spherical homogeneous space $G/N_G(N)$ has the same set of colours as

G/N and, by Proposition 5.4, its wonderful compactification has the same spherical system as that of G/N (see [L01]).

Then we have: $G_x = N$ and

$$G/N \subset \mathbb{P}(V(\lambda_1)) \times \dots \times \mathbb{P}(V(\lambda_s)). \quad (5.3)$$

Define H to be the stabiliser G_v of $v = v_1 + \dots + v_s$ in V . We obtain a spherical homogeneous space, namely G/H , with

$$G/H \subset V. \quad (5.4)$$

Recall that a N -invariant B -eigenvector of $\mathbb{C}(G)$, is uniquely determined by its B -weight, up to a scalar. Moreover, it is (since $\mathbb{C}[G]$ is a unique factorisation domain) a Laurent monomial of irreducible $B \times H$ -eigenvectors of $\mathbb{C}[G]$. Namely, each spherical root can be seen as an integral combination of colours via the map σ' defined above.

Therefore, the B -weights of the H -invariant B -eigenvectors of $\mathbb{C}[G]$ are in correspondence with the given monoid Γ , since it is saturated and $\mathbb{Z}\Gamma$ contains Σ .

The closure X_1 of G/H in V is thus a deformation of X_0 (see Paragraph 2.6).

5.3 The wonderful variety

Once we know the existence of the above wonderful subgroup N , to obtain Theorem 2.3 is enough to show that the closure of G/N in the product of the projective spaces $\mathbb{P}(V(\lambda_i))$ is the wonderful embedding. We shall make use of the following result.

Theorem 5.3 ([Pe05]). *Let G be (connected) semisimple and simply connected, let $\overline{G}$ be its adjoint group $G/Z(G)$. A wonderful $\overline{G}$ -variety X admits a (unique) closed immersion in $\mathbb{P}(V(\lambda))$ if*

- (i) *the dominant weight λ is linear combination with nonzero positive integral coefficients of the colours of X via the map σ in (4.3),*
- (ii) *for any spherical root γ of X , there exists no wonderful $\overline{G}$ -variety X' of rank one with 2γ as spherical root and $S_{X'}^p = S_X^p$.*

Take λ to be equal to $\lambda_1 + \dots + \lambda_s$. The closed immersion of G/N in $\mathbb{P}(V(\lambda_1 + \dots + \lambda_s))$ factorises through the map

$$\phi: \mathbb{P}(V(\lambda_1)) \times \dots \times \mathbb{P}(V(\lambda_s)) \rightarrow \mathbb{P}(V(\lambda_1 + \dots + \lambda_s)), \quad (5.5)$$

the Segre embedding followed by the map induced by the projection of $\otimes_1^s V(\lambda_i)$ onto $V(\lambda_1 + \dots + \lambda_s)$. The closure of G/N in $\prod_1^s \mathbb{P}(V(\lambda_i))$ dominates the closure of G/N in $\mathbb{P}(V(\lambda_1 + \dots + \lambda_s))$. Therefore, if the latter is wonderful then they are equal for the universal property of wonderful varieties.

Since the first condition of the previous theorem is already fulfilled by Proposition 4.4, to conclude that the closure of G/N in $\prod_1^s \mathbb{P}(V(\lambda_i))$ is wonderful we are left to prove the following.

Proposition 5.4. *Let S^p and Σ defined as above from a free and saturated cone $\Gamma \subseteq \Lambda^+$. Then for any $\gamma \in \Sigma$ there exists no wonderful $\overline{G}$ -variety X' of rank one with 2γ as spherical root and $S_{X'}^p = S^p$.*

Proof. The list of possible elements of Σ is in Table 1 and the list of wonderful varieties of rank one is in [W]. The only case of $\gamma \in \Sigma$ such that 2γ is a spherical root is of support of type B_n for $\gamma = \alpha_1 + \dots + \alpha_n$ (where $\alpha_1, \dots, \alpha_n$ are simple roots, namely the simple roots of the support of γ). Let X' be a wonderful variety with $2\alpha_1 + \dots + 2\alpha_n$ as spherical root, then $\alpha_n \in S_{X'}^p$, from Proposition 3.4. Suppose $\alpha_1 + \dots + \alpha_n \in \Sigma$ and $\alpha_n \in S^p$, then there exists a unique j such that $(\lambda_j, \gamma) \neq 0$, it is exactly equal to ω_{α_1} (the fundamental weight associated to α_1). Hence, if we restrict the action of the torus on $(V(\lambda_j)/\mathfrak{g}.v_{\lambda_j})^{G_{v_{\lambda_j}}}$ we should obtain $\alpha_1 + \dots + \alpha_n$ as weight, but this is false and the weight is actually $2\alpha_1 + \dots + 2\alpha_n$. It can be computed by taking the semisimple part of the Levi subgroup G' of G with root subsystem generated by $\alpha_1, \dots, \alpha_n$, with $T' = G' \cap G$ as maximal torus, and considering the T' -weight in $(V(\omega'_1)/\mathfrak{g}'.v')^{G'_{v'}}$ (ω'_1 first fundamental weight of G' with respect to T' , v' highest weight vector in $V(\omega'_1)$). $\square$

Corollary 5.5. *Let $(S^p, \Sigma, \emptyset)$ be a spherical system satisfying the condition of Proposition 5.4, i.e. for any $\gamma \in \Sigma$ there exists no wonderful variety X' of rank one with 2γ as spherical root and $S_{X'}^p = S^p$. Let $\Gamma = \langle \lambda_1, \dots, \lambda_s \rangle_{\mathbb{N}}$ be free and saturated, with S^p equal to the subset of simple roots orthogonal to Γ and $\mathbb{Z}\Gamma \supset \Sigma$, satisfying the conditions of Proposition 4.4, i.e. $\Gamma \subseteq \mathbb{N}\Delta$, via the map σ , and each colour $D \in \Delta$ occurs in the linear combination of at least one weight λ_i .*

Assume moreover that, for a fixed Γ , the set Σ is maximal with the above conditions.

Then $(S^p, \Sigma, \emptyset)$ is the spherical system attached to Γ in the sense of Theorem 4.1 and the dimension of the invariant Hilbert scheme $\text{Hilb}_{\Gamma}^G(V)_0$ equals the rank of Σ .

Notice that the hypotheses of the corollary are fulfilled by $\Gamma = \mathbb{N}\Delta$.

Proof. In the proof of Theorem 2.3 we actually use only the relations between Γ and $(S^p, \Sigma, \emptyset)$ that are here reported as hypotheses of the statement. $\square$

6 Existence of the wonderful subgroup

6.1 Reduction to the primitive cases

In the previous section we remarked that the spherical system arising from a free and saturated cone Γ satisfies the condition of Proposition 5.4. We should now prove that for every spherical system with such a condition there exists a corresponding wonderful subgroup. We want to reduce the proof to some primitive spherical system as in [L01] and [BP]. To do so we use a weaker requirement on the spherical system: we admit all spherical roots which are not simple.

Proposition 6.1. *Let S^p be a subset of simple roots and let Σ be a set of spherical not simple roots such that $(S^p, \Sigma, \emptyset)$ is a spherical system for G of adjoint type. Then there exists a wonderful subgroup N of G such that the wonderful embedding of G/N has the given spherical system.*

In the following we briefly recall some definitions and statements from [L01] which allow us to reduce the proof to finitely many cases. We restrict to the hypotheses of Proposition 6.1.

Let $(S^p, \Sigma, \emptyset)$ be a spherical system, let Δ be its set of colours. A subset $\Delta' \subset \Delta$ is said to be *distinguished* if there exists a linear combination ϕ of the elements of $\rho(\Delta')$, with positive coefficients, such that $\langle \phi, \gamma \rangle \geq 0$ for all $\gamma \in \Sigma$. Let $\Sigma(\Delta')$ denote the maximal subset of spherical roots such that there exists a linear combination ϕ as above such that $\langle \phi, \gamma \rangle > 0$ for all $\gamma \in \Sigma(\Delta')$ and let $V(\Delta')$ denote the subspace spanned by $\rho(\Delta')$ and $\{\gamma^*: \gamma \in \Sigma(\Delta')\}$, where $\{\gamma^*\}$ is the dual basis of Σ .

Let $(S^p, \Sigma, \emptyset)$ be a spherical system and Δ' a distinguished subset of Δ . Let Ξ/Δ' be the subgroup of the weights $\xi \in \Xi$ such that $\langle \rho(D), \xi \rangle = 0$ for all D in Δ' and $\langle \gamma^*, \xi \rangle = 0$ for all γ in $\Sigma(\Delta')$. Let S^p/Δ' be the set $\{\alpha \in S^p: \Delta(\alpha) \subset \Delta'\} \subset S^p$. Let Σ/Δ' be the set of the indecomposable elements of the semigroup $\{\sum_{\gamma \in \Sigma} c_\gamma \gamma \in \Xi/\Delta': c_\gamma \geq 0 \ \forall \gamma \in \Sigma\}$. If the set Σ/Δ' is a basis of Ξ/Δ' , then $(S^p/\Delta', \Sigma/\Delta', \emptyset)$ is a spherical system.

Let $\Phi: X \rightarrow X'$ be a dominant G -morphism between wonderful G -varieties and let Δ_Φ be the set of colours that map dominantly.

Proposition 6.2 ([L01]). *The map $\Phi \mapsto \Delta_\Phi$ is a bijection between the set of dominant G -morphisms with connected fibers of X , onto another wonderful G -variety, and the set of distinguished subsets of Δ_X such that $(S^p/\Delta_\Phi, \Sigma/\Delta_\Phi, \emptyset)$ is a spherical system. The latter is equal to the spherical system of the target of Φ .*

The distinguished subset Δ' of Δ is said to be *smooth* if $V(\Delta')$ is spanned by $\{\gamma^*: \gamma \in \Sigma(\Delta')\}$. In this case $\Sigma/\Delta' = \Sigma \setminus \Sigma(\Delta') \subset \Sigma$. The distinguished subset Δ' of Δ is said to be *parabolic* if $\Sigma(\Delta') = \Sigma$.

Let P be a parabolic subgroup of G and let $L \subset P$ be a Levi subgroup. Let Y be a wonderful L -variety. The *parabolic induction* obtained from Y by P is the wonderful G -variety $G \times_P Y$, where Y is considered as a P -variety where the radical P^r of P acts trivially.

Proposition 6.3 ([L01]). *Let X be a wonderful G -variety and let S' be a subset of S with $(\text{Supp}(\Sigma) \cup S^p) \subset S'$. Then X is obtained by parabolic induction by $P_{S'}^-$, the parabolic subset containing B^- associated to S' . The corresponding G -morphism $\Phi: X \rightarrow G/P_{S'}^-$ is associated to the parabolic distinguished subset $\Delta(S')$.*

A spherical system is said to be *cuspidal* if $\text{Supp}(\Sigma) = S$.

A spherical system $(S^p, \Sigma, \emptyset)$ is said to be *reducible* if there exists a partition of S into two subsets S_1, S_2 such that $S_1 \perp S_2$ and for all γ in Σ $\text{Supp}(\gamma) \subset S_1$ or $\text{Supp}(\gamma) \subset S_2$. In this case $(S^p, \Sigma, \emptyset)$ is the *direct product* of the spherical systems $(S^p \cap S_1, \Sigma_1, \emptyset)$ and $(S^p \cap S_2, \Sigma_2, \emptyset)$, where $\Sigma_i = \{\gamma \in \Sigma: \text{Supp}(\gamma) \subset S_i\}$.

Let $(S^p, \Sigma, \emptyset)$ be a spherical system. Let Δ_1 and Δ_2 be two distinguished subsets of Δ , obviously $\Delta_3 = \Delta_1 \cup \Delta_2$ is distinguished. The subsets Δ_1 and Δ_2 *decompose into fiber product* the spherical system $(S^p, \Sigma, \emptyset)$ if:

- (i) $\Delta_1 \neq \emptyset$, $\Delta_2 \neq \emptyset$ and $\Delta_1 \cap \Delta_2 = \emptyset$,
- (ii) $(S^p/\Delta_i, \Sigma/\Delta_i, \emptyset)$ is a spherical system, for $i = 1, 2, 3$,
- (iii) $(\Sigma \setminus (\Sigma/\Delta_1)) \cap (\Sigma \setminus (\Sigma/\Delta_2)) = \emptyset$,
- (iv) $((S^p/\Delta_1) \setminus S^p) \perp ((S^p/\Delta_2) \setminus S^p)$,
- (v) Δ_1 or Δ_2 is smooth.

Proposition 6.4 ([L01]). *Let $(S^p, \Sigma, \emptyset)$ be a spherical system, let Δ_1 and Δ_2 be two distinguished subsets that decompose $(S^p, \Sigma, \emptyset)$. Let us suppose that, for $i = 1, 2, 3$, there exists a wonderful G -variety X_i with spherical system $(S^p/\Delta_i, \Sigma/\Delta_i, \emptyset)$. Then there exists two G -morphisms $\Phi_1: X_1 \rightarrow X_3$, $\Phi_2: X_2 \rightarrow X_3$ such that the fiber product $X_1 \times_{X_3} X_2$ is a wonderful G -variety with spherical system $(S^p, \Sigma, \emptyset)$.*

The above operation of factorization is a particular case of this decomposition into fiber product. A reducible spherical system, $S = S_1 \sqcup S_2$, is decomposed by $\Delta(S_1)$ and $\Delta(S_2)$, and it corresponds to a direct product of two wonderful varieties.

6.2 Primitive cases

With the above results the proof of Proposition 6.1 is reduced to the spherical systems that are cuspidal and indecomposable. They are called *primitive* spherical systems.

To list the primitive spherical systems we proceed as follows (see [BP]).

Let $(S^p, \Sigma, \emptyset)$ be a spherical system and let Δ be its set of colours. Two spherical roots $\gamma_1, \gamma_2 \in \Sigma$ are said to be *strongly Δ -adjacent* if for all $D \in \Delta(\text{Supp } \gamma_1)$ we have $\langle \rho(D), \gamma_2 \rangle \neq 0$ and, vice versa, for all $D \in \Delta(\text{Supp } \gamma_2)$ we have $\langle \rho(D), \gamma_1 \rangle \neq 0$.

Let Σ' be a subset of Σ . The spherical system $(S^p \cap \text{Supp}(\Sigma'), \Sigma', \emptyset)$ is said to be *strongly Δ -connected* if for every couple of spherical roots $\gamma_1, \gamma_2 \in \Sigma'$ there exists a finite sequence of spherical roots in Σ' , one strongly Δ -adjacent to the next one, the first equal to γ_1 and the last equal to γ_2 . If $\Sigma' \subset \Sigma$ is maximal with this property we say that $(S^p \cap \text{Supp}(\Sigma'), \Sigma', \emptyset)$ is a *strongly Δ -connected component* of $(S^p, \Sigma, \emptyset)$.

Let $(S^p, \Sigma, \emptyset)$ be a spherical system. Let Σ' be a subset of spherical roots. Let $\Delta(\Sigma')$ denote the subset of colours $D \in \Delta(\text{Supp}(\Sigma'))$ such that $\langle \rho(D), \gamma \rangle = 0$ for all $\gamma \in \Sigma \setminus \Sigma'$. We say that $(S^p \cap \text{Supp}(\Sigma'), \Sigma', \emptyset)$ is *erasable* if there exists a nonempty smooth distinguished subset of colours Δ' included in $\Delta(\Sigma')$.

We say that it is *quasi-erasable* if there exists a nonempty distinguished subset of colours Δ' included in $\Delta(\Sigma')$ such that $(S^p/\Delta', \Sigma/\Delta', \emptyset)$ is a spherical system.

Lemma 6.5 ([BP]). *Let $(S^p, \Sigma, \emptyset)$ be a spherical system. Let Σ_1 and Σ_2 be two disjoint subsets of Σ giving two quasi-erasable localizations such that at least one of them is erasable. Then the corresponding subsets $\Delta'_1 \subset \Delta(\Sigma_1)$ and $\Delta'_2 \subset \Delta(\Sigma_2)$ decompose the spherical system $(S^p, \Sigma, \emptyset)$.*

The strongly Δ -connected component $(S^p \cap \text{Supp}(\Sigma'), \Sigma', \emptyset)$ is said to be *isolated* if the partition $\text{Supp}(\Sigma') \sqcup (\text{Supp}(\Sigma) \setminus \text{Supp}(\Sigma'))$ gives a factorization of $(S^p \cap \text{Supp}(\Sigma), \Sigma, \emptyset)$. An isolated component is erasable.

To obtain the list of primitive spherical systems we start from the list of cuspidal strongly Δ -connected spherical systems. Looking at the set of colours of a strongly Δ -connected component we can say whether it is necessarily isolated, erasable, quasi-erasable or none of them. To get all the remaining primitive spherical systems we glue together, in all possible ways, two or more strongly Δ -connected components.

To conclude the proof of Proposition 6.1 we provide here the list of primitive spherical systems with the corresponding wonderful subgroups N . Since the lists for rank one and two are already contained in [W], we restrict to

rank greater than two. The proof that N is wonderful and corresponds to the given spherical system is already given in [L01], [Pe03] and [BP] for types A, C, D. For the other cases the argument is similar.

The unipotent radical N^u of N is equal to the unipotent radical $(N^\circ)^u$ of the connected component N° . A Levi factor K of N is such that $N^\circ = K^\circ N^u$. There exists a parabolic subgroup P with Levi decomposition LP^u with $K^\circ \subseteq L$ and $N^u \subseteq P^u$. We can suppose that P is in standard position. To describe N we provide here the subgroup K up to isogeny and, if N is not reductive, the subset S' of simple roots associated to P and the complementary $\mathfrak{m}$ of the Lie algebra of N^u in the Lie algebra $\mathfrak{n}$ of P^u as representation of the semisimple part $(K^\circ)'$ of K° . Moreover, if necessary, we specify the submodule of $\mathfrak{n}/[\mathfrak{n}, \mathfrak{n}]$ in which $\mathfrak{m}$ is embedded diagonally.

In the following the rank of the connected components of the Dynkin diagram is denoted by n_i , or n if there is only one component.

Type A

- (1) $n \geq 7$ odd, $S^p = \{\alpha_1, \dots, \alpha_{2i-1}, \alpha_{2i+1}, \dots, \alpha_n\}$, $\Sigma = \{\alpha_1 + 2\alpha_2 + \alpha_3, \dots, \alpha_{2i-1} + 2\alpha_{2i} + \alpha_{2i+1}, \dots, \alpha_{n-2} + 2\alpha_{n-1} + \alpha_n\}$. $K = N(Sp_{n+1})$.
- (2) $n_1 = n_2 \geq 3$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha'_1, \dots, \alpha_i + \alpha'_i, \dots, \alpha_{n_1} + \alpha'_{n_2}\}$. $K = N(SL_{n_1+1})$.
- (3) $n = 2p + q$, $p \geq 2$, $q \geq 2$, $S^p = \{\alpha_{p+2}, \dots, \alpha_{p+q-1}\}$, $\Sigma = \{\alpha_1 + \alpha_{p+q+1}, \dots, \alpha_i + \alpha_{p+q+i}, \dots, \alpha_p + \alpha_n; \alpha_{p+1} + \dots + \alpha_{p+q}\}$. $K = S(GL_{p+q} \times GL_{p+1})$.
- (4) $n = 2p + 1$, $p \geq 2$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha_{p+2}, \dots, \alpha_i + \alpha_{p+i+1}, \dots, \alpha_p + \alpha_n; 2\alpha_{p+1}\}$. $K = N(SL_{p+1} \times SL_{p+1})$.
- (5) $n \geq 3$, $S^p = \emptyset$, $\Sigma = \{2\alpha_1, \dots, 2\alpha_i, \dots, 2\alpha_n\}$. $K = N(SO_{n+1})$.
- (6) $n \geq 4$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha_2, \dots, \alpha_i + \alpha_{i+1}, \dots, \alpha_{n-1} + \alpha_n\}$. n even, $S' = S \setminus \{\alpha_n\}$, $K = Sp_n \times GL_1$. n odd, $S' = S \setminus \{\alpha_1, \alpha_n\}$, $K = Sp_{n-1} \times GL_1$, $\mathfrak{m} = V(\omega_1) \subset V(\omega_1)^{\oplus 2}$.

Type B

- (7) $n_1 = n_2 \geq 3$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha'_1, \dots, \alpha_i + \alpha'_i, \dots, \alpha_{n_1} + \alpha'_{n_2}\}$. $K = N(SO_{2n_1+1})$.
- (8) $n \geq 4$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha_2, \dots, \alpha_i + \alpha_{i+1}, \dots, \alpha_{n-1} + \alpha_n\}$. n even, $S' = S \setminus \{\alpha_n\}$, $K = Sp_n \times GL_1$, $\mathfrak{m} = V(\omega_1)$. n odd, $S' = S \setminus \{\alpha_1, \alpha_n\}$, $K = Sp_{n-1} \times GL_1$, $\mathfrak{m} = V(\omega_1) \subset V(\omega_1)^{\oplus 2}$.
- (9) $n = p + q$, $p \geq 2$, $q \geq 1$, $S^p = \{\alpha_{p+2}, \dots, \alpha_n\}$, $\Sigma = \{2\alpha_1, \dots, 2\alpha_i, \dots, 2\alpha_p; 2\alpha_{p+1} + \dots + 2\alpha_n\}$. $K = SO_{p+1} \times SO_{2n-p}$.

(10) $n_1 = n_2 = 2$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha_2, \alpha_2 + \alpha'_2, \alpha'_1 + \alpha'_2\}$. $K = SL_2 \times SL_2 \times SL_2$.

(11) $n = p + q$, $p \geq 3$, $q \geq 1$, $S^p = \alpha_{p+2}, \dots, \alpha_n$, $\Sigma = \{\alpha_1 + \alpha_2, \dots, \alpha_i + \alpha_{i+1}, \dots, \alpha_{p-1} + \alpha_p; 2\alpha_{p+1} + \dots + 2\alpha_n\}$. p even, $S' = S \setminus \{\alpha_p\}$, $K = Sp_p \times N_{SO_{2n-2p+1}}(SO_{2n-2p}) \times GL_1$, $\mathfrak{m} = V(\omega_1)$. p odd, $S' = S \setminus \{\alpha_1, \alpha_p\}$, $K = Sp_{p-1} \times N_{SO_{2n-2p+1}}(SO_{2n-2p}) \times GL_1$, $\mathfrak{m} = V(\omega_1) \subset V(\omega_1)^{\oplus 2}$.

(12) $n \geq 3$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha_2, \dots, \alpha_i + \alpha_{i+1}, \dots, \alpha_{n-1} + \alpha_n; 2\alpha_n\}$. $K = N(GL_n)$.

(13) $n = p + q$, $p \geq 3$, $q \geq 2$, $S^p = \{\alpha_{p+2}, \dots, \alpha_n\}$, $\Sigma = \{\alpha_1 + \alpha_2, \dots, \alpha_i + \alpha_{i+1}, \dots, \alpha_{p-1} + \alpha_p; \alpha_{p+1} + \dots + \alpha_n\}$. p even, $S' = S \setminus \{\alpha_p\}$, $K = Sp_p \times SO_{2n-2p} \times GL_1$, $\mathfrak{m} = V(\omega_1)$. p odd, $S' = S \setminus \{\alpha_1, \alpha_p\}$, $K = Sp_{p-1} \times SO_{2n-2p} \times GL_1$, $\mathfrak{m} = V(\omega_1) \subset V(\omega_1)^{\oplus 2}$.

Type C

(14) $n \geq 6$ even, $S^p = \{\alpha_1, \dots, \alpha_{2i-1}, \alpha_{2i+1}, \dots, \alpha_{n-1}\}$, $\Sigma = \{\alpha_1 + 2\alpha_2 + \alpha_3, \dots, \alpha_{2i-1} + 2\alpha_{2i} + \alpha_{2i+1}, \dots, \alpha_{n-3} + 2\alpha_{n-2} + \alpha_{n-1}; 2\alpha_{n-1} + 2\alpha_n\}$. $K = N(Sp_n \times Sp_n)$.

(15) $n_1 = n_2 \geq 3$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha'_1, \dots, \alpha_i + \alpha'_i, \dots, \alpha_{n_1} + \alpha'_{n_2}\}$. $K = N(Sp_{2n})$.

(16) $n \geq 3$, $S^p = \emptyset$, $\Sigma = \{2\alpha_1, \dots, 2\alpha_i, \dots, 2\alpha_n\}$. $K = N(GL_n)$.

(17) $n = p + q$, $p \geq 4$, $q \geq 2$, $S^p = \{\alpha_1, \dots, \alpha_{2i-1}, \alpha_{2i+1}, \dots, \alpha_{p+1}; \alpha_{p+3}, \dots, \alpha_n\}$, $\Sigma = \{\alpha_1 + 2\alpha_2 + \alpha_3, \dots, \alpha_{2i-1} + 2\alpha_{2i} + \alpha_{2i+1}, \dots, \alpha_{p-1} + 2\alpha_p + \alpha_{p+1}; \alpha_{p+1} + 2\alpha_{p+2} + \dots + 2\alpha_{n-1} + \alpha_n\}$. $K = Sp_{2p+2} \times Sp_{2n-2p-2}$.

(18) $n = p + q$, $p \geq 4$, $q \geq 1$, $S^p = \{\alpha_3, \dots, \alpha_{p-2}; \alpha_{p+2}, \dots, \alpha_n\}$, $\Sigma = \{\alpha_1 + \alpha_p, \alpha_2 + \dots + \alpha_{p-1}, \alpha_{p+1} + 2\alpha_{p+2} + \dots + 2\alpha_{n-1} + \alpha_n\}$. $S' = S \setminus \{\alpha_{p-1}\}$, $K = SL_p \times Sp_{2n-2p-2} \times GL_1$, $\mathfrak{m} = V(\omega'_1)$.

(19) $n_1, n_2 \geq 3$, $S^p = \{\alpha_3, \dots, \alpha_{n_1}; \alpha'_3, \dots, \alpha'_{n_2}\}$, $\Sigma = \{\alpha_1 + \alpha'_1, \alpha_1 + 2\alpha_2 + \dots + 2\alpha_{n_1-1} + \alpha_{n_1}, \alpha'_1 + 2\alpha'_2 + \dots + 2\alpha'_{n_2-1} + \alpha'_{n_2}\}$. $K = SL_2 \times Sp_{2n_1-2} \times Sp_{2n_2-2}$.

(20) $n = p + q$, $p \geq 3$, $q \geq 1$, $S^p = \{\alpha_{p+2}, \dots, \alpha_n\}$, $\Sigma = \{\alpha_1 + \alpha_2, \dots, \alpha_i + \alpha_{i+1}, \dots, \alpha_{p-1} + \alpha_p; \alpha_p + 2\alpha_{p+1} + \dots + 2\alpha_{n-1} + \alpha_n\}$. p even, $S' = S \setminus \{\alpha_1\}$, $K = Sp_p \times Sp_{2n-p-2} \times GL_1$, $\mathfrak{m} = V(\omega_1)$. p odd, $S' = S \setminus \{\alpha_1\}$, $K = Sp_{p-1} \times Sp_{2n-p-1} \times GL_1$, $\mathfrak{m} = V(\omega'_1)$.

Type D

(21) $n \geq 6$ even, $S^p = \{\alpha_1, \dots, \alpha_{2i-1}, \alpha_{2i+1}, \dots, \alpha_{n-1}\}$, $\Sigma = \{\alpha_1 + 2\alpha_2 + \alpha_3, \dots, \alpha_{2i-1} + 2\alpha_{2i} + \alpha_{2i+1}, \dots, \alpha_{n-3} + 2\alpha_{n-2} + \alpha_{n-1}; 2\alpha_n\}$. $K = N(GL_n)$.

(22) $n = p + q$, $p \geq 2$, $q \geq 2$, $S^p = \{\alpha_{p+2}, \dots, \alpha_n\}$, $\Sigma = \{2\alpha_1, \dots, 2\alpha_i, \dots, 2\alpha_p; 2\alpha_{p+1} + \dots + 2\alpha_{n-2} + \alpha_{n-1} + \alpha_n\}$. $K = S(O_{p+1} \times O_{2n-p-1})$.

- (23) $n \geq 7$ odd, $S^p = \{\alpha_1, \dots, \alpha_{2i-1}, \alpha_{2i+1}, \dots, \alpha_{n-2}\}$, $\Sigma = \{\alpha_1 + 2\alpha_2 + \alpha_3, \dots, \alpha_{2i-1} + 2\alpha_{2i} + \alpha_{2i+1}, \dots, \alpha_{n-4} + 2\alpha_{n-3} + \alpha_{n-2}; \alpha_{n-1} + \alpha_{n-2} + \alpha_n\}$. $K = GL_n$.
- (24) $n_1 = n_2 \geq 4$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha'_1, \dots, \alpha_i + \alpha'_i, \dots, \alpha_{n_1} + \alpha'_{n_2}\}$. $K = N(SO_{2n_1})$.
- (25) $n \geq 4$, $S^p = \emptyset$, $\Sigma = \{2\alpha_1, \dots, 2\alpha_i, \dots, 2\alpha_n\}$. $K = N(SO_n \times SO_n)$.
- (26) $n = 4$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha_2 + \alpha_3, \alpha_3 + \alpha_2 + \alpha_4, \alpha_4 + \alpha_2 + \alpha_1\}$. $K = N(G_2)$.
- (27) $n \geq 4$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha_2, \dots, \alpha_i + \alpha_{i+1}, \dots, \alpha_{n-2} + \alpha_n; \alpha_{n-2} + \alpha_n\}$. n even, $S' = S \setminus \{\alpha_1, \alpha_{n-1}, \alpha_n\}$, $K = Sp_{n-2} \times GL_1$, $\mathfrak{m} = V(\omega_1) \subset V(\omega_1)^{\oplus 3}$. n odd, $S' = S \setminus \{\alpha_{n-1}, \alpha_n\}$, $K = Sp_{n-1} \times GL_1$, $\mathfrak{m} = V(\omega_1) \subset V(\omega_1)^{\oplus 2}$.
- (28) $n = p + q$, $p \geq 3$, $q \geq 2$, $S^p = \{\alpha_{p+2}, \dots, \alpha_n\}$, $\Sigma = \{\alpha_1 + \alpha_2, \dots, \alpha_i + \alpha_{i+1}, \dots, \alpha_{n-2} + \alpha_n; 2\alpha_{p+1} + \dots + 2\alpha_{n-2} + \alpha_{n-1} + \alpha_n\}$. p odd, $S' = S \setminus \{\alpha_1, \alpha_p\}$, $K = Sp_{p-1} \times SO_{2q-1} \times GL_1$, $\mathfrak{m} = V(\omega_1) \subset V(\omega_1)^{\oplus 2}$. p even, $S' = S \setminus \{\alpha_p\}$, $K = Sp_p \times SO_{2q-1} \times GL_1$, $\mathfrak{m} = V(\omega_1)$.

Type E

- (29) $n = 7$ (or 8), $S^p = \{\alpha_2, \alpha_3, \alpha_4, \alpha_5\}$, $\Sigma = \{2\alpha_1 + 2\alpha_3 + 2\alpha_4 + \alpha_2 + \alpha_5, 2\alpha_6 + 2\alpha_5 + 2\alpha_4 + \alpha_2 + \alpha_3, 2\alpha_7, (2\alpha_8)\}$. $K = N(E_6 \times GL_1), E_7 \times SL_2$.
- (30) $n = 7$, $S^p = \{\alpha_2, \alpha_5, \alpha_7\}$, $\Sigma = \{2\alpha_1, 2\alpha_3, \alpha_2 + 2\alpha_4 + \alpha_5, \alpha_5 + 2\alpha_6 + \alpha_7\}$. $K = Spin_{12} \times SL_2$.
- (31) $n_1 = n_2 = 6, 7, 8$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha'_1, \dots, \alpha_i + \alpha'_i, \dots, \alpha_{n_1} + \alpha'_{n_2}\}$. $K = N(E_{n_1})$.
- (32) $n = 6$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha_6, \alpha_3 + \alpha_5, 2\alpha_2, 2\alpha_4\}$. $K = SL_6 \times SL_2$.
- (33) $n = 6, 7, 8$, $S^p = \emptyset$, $\Sigma = \{2\alpha_1, \dots, 2\alpha_i, \dots, 2\alpha_n\}$. $K = Sp_8, SL_8, Spin_{16}$.
- (34) $n = 6, 7, 8$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha_3, \alpha_2 + \alpha_4, \alpha_3 + \alpha_4, \dots, \alpha_i + \alpha_{i+1}, \dots, \alpha_{n-1} + \alpha_n\}$. $n = 6$, $S' = S \setminus \{\alpha_2\}$, $K = Sp_6 \times GL_1$, $\mathfrak{m} = V(\omega_1)$. $n = 7$, $S' = S \setminus \{\alpha_2, \alpha_7\}$, $K = Sp_6 \times GL_1$, $\mathfrak{m} = V(\omega_1) \subset V(\omega_1)^{\oplus 2}$. $n = 8$, $S' = S \setminus \{\alpha_2\}$, $K = Sp_8 \times GL_1$, $\mathfrak{m} = V(\omega_1)$.
- (35) $n = 8$, $S^p = \{\alpha_2, \alpha_3, \alpha_4, \alpha_5\}$, $\Sigma = \{2\alpha_1 + 2\alpha_3 + 2\alpha_4 + \alpha_2 + \alpha_5, 2\alpha_6 + 2\alpha_5 + 2\alpha_4 + \alpha_2 + \alpha_3, \alpha_7 + \alpha_8\}$. $S' = S \setminus \{\alpha_7\}$, $K = F_4 \times SL_2 \times GL_1$, $\mathfrak{m} = V(\omega'_1)$.
- (36) $n = 6$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha_6, \alpha_3 + \alpha_5, \alpha_2 + \alpha_4\}$. $S' = S \setminus \{\alpha_4\}$, $K = N_{SL_3 \times SL_3}(SL_3) \times SL_2 \times GL_1$, $\mathfrak{m} = V(\omega'_1)$.
- (37) $n = 7$, $S^p = \{\alpha_2, \alpha_5, \alpha_7\}$, $\Sigma = \{\alpha_1 + \alpha_3, \alpha_2 + 2\alpha_4 + \alpha_5, \alpha_5 + 2\alpha_6 + \alpha_7\}$. $S' = S \setminus \{\alpha_3\}$, $K = SL_2 \times Sp_6 \times GL_1$, $\mathfrak{m} = V(\omega_1)$.

Type F

- (38) $n_1 = n_2 = 4$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha'_1, \alpha_2 + \alpha'_2, \alpha_3 + \alpha'_3, \alpha_4 + \alpha'_4\}$. $K = F_4$.
- (39) $n = 4$, $S^p = \emptyset$, $\Sigma = \{2\alpha_1, 2\alpha_2, 2\alpha_3, 2\alpha_4\}$. $K = Sp_6 \times SL_2$.

(40) $n = 4$, $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha_2, 2\alpha_3, 2\alpha_4\}$. $S' = S \setminus \{\alpha_2\}$, $K = SL_2 \times SO_3 \times GL_1$, $\mathfrak{m} = V(\omega_1)$.

In type G there are no primitive spherical systems with rank greater than two.

References

- [A] Ahiezer, D.N., *Equivariant completions of homogeneous algebraic varieties by homogeneous divisors*, Ann. Global Anal. Geom. **1** (1983), no. 1, 49–78.
- [AB] Alexeev, V. and Brion, M., *Moduli of affine schemes with reductive group action*, J. Algebraic Geom., **14** (2005), no. 1, 83–117.
- [BK] Borho, W. and Kraft, H., *Über Bahnen und deren Deformationen bei linearen Aktionen reduktiver Gruppen*, Comment. Math. Helv., **54** (1979), 61–104.
- [Bra] Bravi, P., *Wonderful varieties of type E*, preprint.
- [BP] Bravi, P. and Pezzini, G., *Wonderful varieties of type D*, Represent. Theory **9** (2005), 578–637.
- [Bri86] Brion, M., *Quelques propriétés des espaces homogènes sphériques*, Manuscripta Math., **55** (1986), 191–198.
- [Bri89] Brion, M., *On spherical varieties of rank one (after D. Ahiezer, A. Huckleberry, D. Snow)*, Group actions and invariant theory (Montreal, PQ, 1988), CMS Conf. Proc., 10, Amer. Math. Soc., Providence, RI, 1989, 31–41.
- [F] Foschi, A., *Variétés magnifiques et polytopes moment*, Ph.D. Thesis, Institut Fourier, Université J. Fourier, Grenoble, 1998.
- [HS] Huckleberry, A., and Snow, D., *Almost-homogeneous Kähler manifolds with hypersurface orbits*, Osaka J. Math. **19** (1982), no. 4, 582–593.
- [J] Jansou, S., *Déformations invariantes des cônes de vecteurs primitifs*, arXiv:math.AG/0506133 v1 .
- [K90] Knop, F., *Weylgruppe und Momentabbildung*, Invent. Math. **99** (1990), 1–23.

- [K96] Knop, F., *Automorphisms, root systems, and compactifications of homogeneous varieties*, J. Amer. Math. Soc. **9** (1996), no. 1, 153–174.
- [L96] Luna, D., *Toute variété magnifique est sphérique*, Transform. Groups **1** (1996), no. 3, 249–258.
- [L97] Luna, D., *Grosses cellules pour les variétés sphériques*, Algebraic groups and Lie groups, Austral. Math. Soc. Lect. Ser., 9, Cambridge Univ. Press, Cambridge, 1997, 267–280.
- [L01] Luna, D., *Variétés sphériques de type A*, Inst. Hautes Études Sci. Publ. Math. **94** (2001), 161–226.
- [L05] Luna, D., *Espaces homogènes modèles*, Seminar talk, Institut Fourier, Grenoble, January 2005.
- [Pa] Panyushev, D., *On deformation method in invariant theory*, Ann. Inst. Fourier **47** (1997), no. 4, 985–1012.
- [Pe03] Pezzini, G., *Wonderful varieties of type C*, Ph.D. Thesis, Dipartimento di Matematica G. Castelnuovo, Università La Sapienza, Roma, 2003.
- [Pe05] Pezzini, G., *Simple immersions of wonderful varieties*, arXiv:math.RT/0506601 .
- [VP] Vinberg, E.B. and Popov, V.L., *On a class of quasihomogeneous affine varieties*, Math. USSR Izv. **6** (1972), 743–758.
- [W] Wasserman, B., *Wonderful varieties of rank two*, Transform. Groups **1** (1996), no. 4, 375–403.

Dipartimento di Matematica, Università di Padova, via G.B. Belzoni 7, 35131 Padova, Italy
 bravi@math.unipd.it

Mathematisches Institut, Universität zu Köln, Weyertal Str. 86-90, 50931 Köln, Germany
 scupit@mi.uni-koeln.de